

セジロウカ、トビロウカ、ヒメトビウカ、コブメイガの誘殺状況(平成28年)

月・日	セジロウカ					トビロウカ					ヒメトビウカ					コブメイガ		
	諫早		佐世保	北有馬	岐宿	諫早		佐世保	北有馬	岐宿	諫早		佐世保	北有馬	岐宿	諫早	生息量(頭/10m)	フェロモントラップ
	白熱	ネット	白熱	白熱	白熱	白熱	ネット	白熱	白熱	白熱	白熱	ネット	白熱	白熱	白熱			
5/1	0	0				0	0				0	0						
5/2	0	0				0	0				0	0						
5/3	0	1				0	0				0	0						
5/4	0	1				0	0				0	0						
5/5	0	0				0	0				0	0						
5/6	0	0				0	0				0	0						
5/7	0	0				0	0				0	0						
5/8	0	0				0	0				0	0						
5/9	0	0				0	0				0	0						
5/10	0	0				0	0				0	0						
5/11	0	0				0	0				0	0						
5/12	0	0				0	0				0	0						
5/13	0	0				0	0				0	0						
5/14	0	0				0	0				0	0						
5/15	0	0				0	0				0	0						
5/16	0	0				0	0				0	0						
5/17	0	0				0	0				0	0						
5/18	0	0				0	0				0	0						
5/19	0	0				0	0				0	0						
5/20	0	0				0	0				0	0						
5/21	0	0				0	0				0	0						
5/22	0	0				0	0				0	0						
5/23	0	0				0	0				0	0						
5/24	0	0				0	0				0	0						
5/25	0	0				0	0				0	0					0	
5/26	0	0				0	0				0	0				0	0	
5/27	0	0				0	0				0	0				0	0	
5/28	0	0				0	0				0	0				0	0	
5/29	0	0				0	0				0	0				0	0	
5/30	0	0				0	0				0	2				0	0	
5/31	0	0				0	0				0	0				0	0	

月・日	セジロウカ					トビロウカ					ヒメトビウカ					コブメイガ	
	諫早		佐世保	北有馬	岐宿	諫早		佐世保	北有馬	岐宿	諫早		佐世保	北有馬	岐宿	フェロモントラップA	フェロモントラップB
	白熱	ネット	白熱	白熱	白熱	白熱	ネット	白熱	白熱	白熱	白熱	ネット	白熱	白熱	白熱		
6/1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
6/2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
6/3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
6/4	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0
6/5	4	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0
6/6	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
6/7	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
6/8	3	0	0	0	0	1	0	0	1	0	0	0	0	0	0	0	0
6/9	1	0	0	0	0	2	0	0	0	0	0	0	0	0	0	1	0
6/10	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
6/11	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
6/12	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
6/13	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
6/14	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
6/15	20	0	6	0	2	0	0	0	0	0	0	0	0	0	0	0	0
6/16	116	0	0	0	0	0	0	0	0	0	0	0	0	0	2	0	0
6/17	10	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
6/18	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
6/19	22	5	1	0	0	2	0	0	0	0	0	0	0	0	0	0	0
6/20	2	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
6/21	65	3	1	52	3	0	0	0	0	0	0	0	0	0	1	0	0
6/22	34	0	0	3	0	2	0	0	0	0	0	0	0	0	7	0.5	0
6/23	71	0	0	1	0	1	0	0	0	0	1	0	0	0	10	0.5	0
6/24	295	-	4	0	0	1	-	0	0	0	2	-	0	0	35	1	0
6/25	3	-	0	0	0	0	-	0	0	0	0	-	0	0	8	1	0
6/26	0	-	3	0	0	0	-	0	0	0	0	-	0	0	50	12.5	0
6/27	14	-	0	1	1	1	-	0	0	0	0	-	0	0	6	5	0
6/28	56	-	0	0	2	0	-	0	0	0	0	-	0	0	3	1	0
6/29	58	-	0	2	1	0	-	0	0	0	3	-	0	0	4	0.5	0
6/30	7	-	0	0	0	0	-	0	0	0	0	-	0	0	3	0.5	0

白熱:60W白熱灯(予察灯)、ネット:ネットトラップ
 フェロモントラップA(4/20植え早期水稻に1個設置)、フェロモントラップB(6/3植え普通期水稻に2個設置の平均値)
 データ欄の空欄は未調査、-は欠測

月・日	セジロウカ					ヒイロウカ					ヒメヒロウカ					コブメカ	
	諫早		佐世保	北有馬	岐宿	諫早		佐世保	北有馬	岐宿	諫早		佐世保	北有馬	岐宿	諫早	
	白熱	ネット	白熱	白熱	白熱	白熱	ネット	白熱	白熱	白熱	白熱	ネット	白熱	白熱	白熱	フェロモントラップA	フェロモントラップB
7/1	1	-	0	0	0	0	-	0	0	0	0	-	0	0	0	0	0
7/2	2	-	0	0	1	0	-	0	0	0	0	-	0	0	0	2	0
7/3	0	-	0	0	0	0	-	0	0	0	0	-	0	0	0	2	0.5
7/4	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0.5
7/5	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	0
7/6	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	5	6
7/7	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	3	1
7/8	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.5
7/9	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
7/10	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	5	0.5
7/11	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.5
7/12	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0
7/13	27	1	0	0	4	2	0	0	0	0	0	0	0	0	0	0	0
7/14	71	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2.5
7/15	6	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1.5
7/16	153	0	0	0	8	1	0	0	0	0	3	0	0	0	0	1	6
7/17	50	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	3
7/18	41	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	3.5
7/19	72	0	0	0	4	0	0	0	0	0	4	1	0	0	0	0	4.5
7/20	7	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
7/21	6	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0	2
7/22	6	0	0	0	1	0	0	0	0	0	2	0	0	0	0	0	0.5
7/23	15	0	0	0	0	0	0	0	0	0	4	0	0	0	0	0	0.5
7/24	12	1	0	0	1	0	0	0	0	0	3	0	0	0	0	0	0
7/25	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	4
7/26	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	7
7/27	5	0	0	0	0	0	0	0	0	0	0	0	0	0	0	3	0.5
7/28	4	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	5.5
7/29	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	7
7/30	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	3	8
7/31	14	0	0	0	0	0	0	0	0	0	1	0	0	0	0	3	8.5

月・日	セジロウカ					ヒイロウカ					ヒメヒロウカ					コブメカ	
	諫早		佐世保	北有馬	岐宿	諫早		佐世保	北有馬	岐宿	諫早		佐世保	北有馬	岐宿	諫早	
	白熱	ネット	白熱	白熱	白熱	白熱	ネット	白熱	白熱	白熱	白熱	ネット	白熱	白熱	白熱	フェロモントラップA	フェロモントラップB
8/1	8	0	0	0	1	0	0	0	0	0	0	0	0	0	0	2	8.5
8/2	12	1	0	0	1	0	0	0	0	0	0	0	0	0	0	6	6
8/3	3	0	1	0	1	0	0	0	0	0	0	0	0	0	0	5	4
8/4	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	5	7.5
8/5	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	5.5
8/6	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	6
8/7	3	0	1	0	1	0	0	0	0	0	0	0	0	0	0	4	1.5
8/8	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	5
8/9	4	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	2
8/10	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
8/11	5	1	0	1	0	0	0	0	0	0	1	0	0	0	0	1	1.5
8/12	11	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1
8/13	17	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0
8/14	27	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	1
8/15	10	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
8/16	6	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0
8/17	3	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0
8/18	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.5
8/19	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	0.5
8/20	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0
8/21	2	0	0	0	0	0	0	0	0	0	1	0	0	0	0	2	0.5
8/22	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	0.5
8/23	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	2
8/24	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
8/25	4	0	0	0	0	0	0	0	0	0	2	0	0	0	0	2	4.5
8/26	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1	5.5
8/27	7	0	3	0	0	0	0	0	0	0	2	0	0	0	0	2	9
8/28	2	18	0	0	0	0	0	0	0	0	0	0	0	0	0	14	16
8/29	2	2	0	3	0	0	0	0	0	0	0	1	0	0	0	5	3.5
8/30	4	29	0	1	1	0	0	0	0	0	1	7	0	2	0	5	24
8/31	7	13	0	4	6	0	0	0	0	0	0	0	0	4	0	0	6.5

白熱:60W白熱灯(予察灯)、ネット:ネットトラップ
フェロモントラップA(4/20植え早期水稲に1個設置、7/13より6/15植え普通期水稲に移動)、フェロモントラップB(6/3植え普通期水稲に2個設置の平均値
データ欄の空欄は未調査、-は欠測

月・日	セイロウカ					トイロウカ					ヒメトウカ					コノメカ	
	諫早		佐世保	北有馬	岐宿	諫早		佐世保	北有馬	岐宿	諫早		佐世保	北有馬	岐宿	諫早	
	白熱	ネット	白熱	白熱	白熱	白熱	ネット	白熱	白熱	白熱	白熱	ネット	白熱	白熱	白熱	フェロモントラップA	フェロモントラップB
9/1	10	0	0	0	2	0	0	0	0	0	2	0	0	0	1	10	6
9/2	10	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	8
9/3	2	-	0	0	0	0	-	0	0	0	-	0	0	0	7	5.5	
9/4	2	-	0	0	4	0	-	0	0	0	2	-	0	0	8	4	
9/5	6	0	0	0	0	0	0	0	0	0	1	0	0	0	19	17	
9/6	11	1	1	0	0	0	0	0	0	0	3	1	0	0	15	4.5	
9/7	0	8	1	0	1	0	2	0	0	0	0	3	0	0	14	8.5	
9/8	7	2	0	0	0	1	0	0	0	0	0	1	0	0	3	15	
9/9	6	2	0	0	0	3	1	0	13	0	1	1	0	0	2	12.5	
9/10	5	0	0	5	0	0	0	14	0	0	0	0	0	0	0	5	
9/11	4	3	0	8	5	1	0	32	3	0	0	0	0	1	14	1	
9/12	6	0	0	36	0	3	0	5	0	1	0	0	2	0	1	2	
9/13	7	0	0	3	0	0	0	1	0	0	0	0	0	0	1	4.5	
9/14	28	0	0	0	1	8	0	0	1	1	0	0	0	0	5	2	
9/15	18	0	0	9	1	0	0	0	0	0	0	0	0	0	1	2	
9/16	2	終了	1	0	0	1	終了	0	0	1	0	終了	0	0	終了	終了	
9/17	1		0	0	0	0		0	0	0	0		0	0			
9/18	5		0	0	1	0		0	0	1	0		0	0			
9/19	1		0	0	0	0		0	0	0	0		0	0			
9/20	0		0	0	0	0		0	0	0	0		0	0			
9/21	0		0	0	0	0		0	0	1	0		0	0			
9/22	0		0	0	0	2		0	0	0	0		0	0			
9/23	1		0	0	0	1		0	0	0	1		0	0			
9/24	14		0	0	1	11		0	0	1	0		0	0			
9/25	2		0	0	3	3		0	0	7	1		0	0			
9/26	6		0	0	0	6		0	0	0	0		0	0			
9/27	0		0	0	0	1		0	0	0	0		0	0			
9/28	7		0	0	1	0		0	0	0	0		0	0			
9/29	5		0	0	0	0		0	0	0	1		0	0			
9/30	2		0	0	0	1		0	0	0	0		0	0			

白熱: 60W白熱灯(予察灯)、ネット: ネットトラップ
フェロモントラップA(4/20植え早期水稲に1個設置、7/13より6/15植え普通期水稲に移動)、フェロモントラップB(6/3植え普通期水稲に2個設置の平均値
データ欄の空欄は未調査、- は欠測