

課題名	9. みかん灰色かび病の防除適期について																																																																																																		
成果の要約	2回散布区では落弁期・生理落果期散布区の防除効果が最も高かった。																																																																																																		
成	<p>1. 3回散布の防除効果が最も高かった。</p> <p>2. 2回散布では落弁期・生理落果期散布区の防除効果が最も高かった。</p> <p>3. 1回散布区では昭和56年～57年は生理落果期の防除効果が高かったが、昭和58年、59年、60年は落弁期が高かった。</p> <p>第1表 みかん灰色かび病の防除適期</p>																																																																																																		
績	<table border="1"> <thead> <tr> <th rowspan="2">区</th> <th colspan="3">薬剤散布時期</th> <th colspan="5">調査年度</th> </tr> <tr> <th>満開期</th> <th>落弁期</th> <th>生理落果期</th> <th>昭和56年</th> <th>昭和57年</th> <th>昭和58年</th> <th>昭和59年</th> <th>昭和60年</th> </tr> </thead> <tbody> <tr> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>(%)</td> <td></td> <td></td> </tr> <tr> <td>1</td> <td>○</td> <td>○</td> <td>○</td> <td>7.2</td> <td>3.7</td> <td>2.6</td> <td>1.0</td> <td>3.7</td> </tr> <tr> <td>2</td> <td>○</td> <td>○</td> <td>—</td> <td>13.5</td> <td>10.0</td> <td>4.0</td> <td>4.0</td> <td>7.5</td> </tr> <tr> <td>3</td> <td>—</td> <td>○</td> <td>○</td> <td>8.0</td> <td>7.0</td> <td>3.0</td> <td>2.5</td> <td>6.5</td> </tr> <tr> <td>4</td> <td>○</td> <td>—</td> <td>○</td> <td>10.0</td> <td>9.0</td> <td>4.0</td> <td>3.5</td> <td>7.5</td> </tr> <tr> <td>5</td> <td>○</td> <td>—</td> <td>—</td> <td>18.0</td> <td>21.5</td> <td>9.1</td> <td>9.0</td> <td>12.3</td> </tr> <tr> <td>6</td> <td>—</td> <td>○</td> <td>—</td> <td>11.0</td> <td>13.0</td> <td>7.0</td> <td>6.0</td> <td>8.0</td> </tr> <tr> <td>7</td> <td>—</td> <td>—</td> <td>○</td> <td>9.0</td> <td>10.0</td> <td>7.5</td> <td>8.0</td> <td>9.5</td> </tr> <tr> <td>8</td> <td>—</td> <td>—</td> <td>—</td> <td>31.0</td> <td>32.0</td> <td>12.0</td> <td>11.5</td> <td>14.0</td> </tr> </tbody> </table>	区	薬剤散布時期			調査年度					満開期	落弁期	生理落果期	昭和56年	昭和57年	昭和58年	昭和59年	昭和60年							(%)			1	○	○	○	7.2	3.7	2.6	1.0	3.7	2	○	○	—	13.5	10.0	4.0	4.0	7.5	3	—	○	○	8.0	7.0	3.0	2.5	6.5	4	○	—	○	10.0	9.0	4.0	3.5	7.5	5	○	—	—	18.0	21.5	9.1	9.0	12.3	6	—	○	—	11.0	13.0	7.0	6.0	8.0	7	—	—	○	9.0	10.0	7.5	8.0	9.5	8	—	—	—	31.0	32.0	12.0	11.5	14.0
区	薬剤散布時期			調査年度																																																																																															
	満開期	落弁期	生理落果期	昭和56年	昭和57年	昭和58年	昭和59年	昭和60年																																																																																											
						(%)																																																																																													
1	○	○	○	7.2	3.7	2.6	1.0	3.7																																																																																											
2	○	○	—	13.5	10.0	4.0	4.0	7.5																																																																																											
3	—	○	○	8.0	7.0	3.0	2.5	6.5																																																																																											
4	○	—	○	10.0	9.0	4.0	3.5	7.5																																																																																											
5	○	—	—	18.0	21.5	9.1	9.0	12.3																																																																																											
6	—	○	—	11.0	13.0	7.0	6.0	8.0																																																																																											
7	—	—	○	9.0	10.0	7.5	8.0	9.5																																																																																											
8	—	—	—	31.0	32.0	12.0	11.5	14.0																																																																																											
要	<p>注) 表中の数字は発病果率</p> <p>(S 60長崎果試)</p>																																																																																																		
普及上の留意点	昭和61年病害虫防除基準の記載事項を補強する成果である。																																																																																																		